



Direct lysis buffer for PCR/RPA/LAMP (for DNA&RNA)

✉ info@ezassay.com

🌐 www.ezassay.com

EZassay Biotechnology Ltd.

Catalog Code: LYS-RD-100
LYS-RD-1000

CONTENTS

<u>Contents</u>	<u>Page</u>
Brief introduction	1
Materials supplied	1
Procedure(example is for reference only)	1

Brief introduction

This kit uses a specially formulated lysis system that can quickly release nucleic acids from samples, especially for DNA viruses, RNA viruses, prokaryotic bacteria (including gram-negative bacteria and gram-positive bacteria) and other samples. It enables amplifications with samples without extraction and purification such as PCR, RPA, and LAMP. It saves time, labour and cost. This is very important of point-of-care test.

Materials supplied

Item	LYS-RD-100	LYS-RD-1000	Store
Direct lysis buffer for PCR/RPA/LAMP	5ml *1	5ml *10	2-8°C
RNase inhibitor	50µL *1	50µL *10	-20°C

Procedure(example is for reference only)

1. Preparation. Equilibrate the reagents to room temperature prior to use. Depending on the number of samples, a mixture of direct lysis buffer and RNase inhibitor is freshly prepared.

Direct lysis buffer	RNase inhibitor	Ratio
50µl	0.5µl	100:1

2. Harvest cells. Take 200 µl of the sample to be tested and add it to a 1.5 ml centrifuge tube and centrifuge at 12,000 rpm for 10 minutes.

3. Direct lysis. Remove the supernatant, add 50 µl of direct lysis buffer (RNase inhibitor added) to the pellet. Mix well by vortex. Centrifuge at 12,000 rpm for 1minute, and take 1~5 µl of supernatant for amplification such as PCR, RPA, and LAMP.